

# The EUV Emission-Line Spectrum of OY Carinae in Superoutburst: Scattering in the Wind

*C. W. Mauche, J. C. Raymond*

This article was submitted to  
Astro Physical Plasmas: Codes, Models and Observation  
Mexico City, Mexico  
October 25-29, 1999

**U.S. Department of Energy**

Lawrence  
Livermore  
National  
Laboratory

**November 21, 1999**

## DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

This work was performed under the auspices of the United States Department of Energy by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.

This report has been reproduced directly from the best available copy.

Available electronically at <http://www.doc.gov/bridge>  
Available for a processing fee to U.S. Department of Energy  
And its contractors in paper from  
U.S. Department of Energy  
Office of Scientific and Technical Information  
P.O. Box 62  
Oak Ridge, TN 37831-0062  
Telephone: (865) 576-8401  
Facsimile: (865) 576-5728  
E-mail: [reports@adonis.osti.gov](mailto:reports@adonis.osti.gov)

Available for the sale to the public from  
U.S. Department of Commerce  
National Technical Information Service  
5285 Port Royal Road  
Springfield, VA 22161  
Telephone: (800) 553-6847  
Facsimile: (703) 605-6900  
E-mail: [orders@ntis.fedworld.gov](mailto:orders@ntis.fedworld.gov)  
Online ordering: <http://www.ntis.gov/ordering.htm>  
Or  
Lawrence Livermore National Laboratory  
Technical Information Department's Digital Library  
<http://www.llnl.gov/tid/Library.html>

# THE EUV EMISSION-LINE SPECTRUM OF OY CARINAE IN SUPEROUTBURST: SCATTERING IN THE WIND

Christopher W. Mauche

Lawrence Livermore National Laboratory, Livermore, CA

and

John C. Raymond

Harvard-Smithsonian Center for Astrophysics, Cambridge, MA

## RESUMEN

The *Extreme Ultraviolet Explorer* satellite observed the cataclysmic variable OY Carinae in superoutburst for 3 days in 1997 March. The resulting 80–180 Å spectrum, which contains broad (FWHM  $\sim 2000$  km s $^{-1}$ ) emission lines of O V–VI, Ne V–VI, Mg V–VI, and Fe VI–VIII, is well modeled by scattering of boundary layer radiation in the system’s accretion disk wind.

## ABSTRACT

The *Extreme Ultraviolet Explorer* satellite observed the cataclysmic variable OY Carinae in superoutburst for 3 days in 1997 March. The resulting 80–180 Å spectrum, which contains broad (FWHM  $\sim 2000$  km s $^{-1}$ ) emission lines of O V–VI, Ne V–VI, Mg V–VI, and Fe VI–VIII, is well modeled by scattering of boundary layer radiation in the system’s accretion disk wind.

**Key Words:** BINARIES: ECLIPSING — NOVAE, CATAclysmic VARIABLES — STARS: INDIVIDUAL (OY CARINAE) — STARS: WINDS, OUTFLOWS — ULTRAVIOLET: STARS

Cataclysmic variables (CVs) are a diverse class of semidetached binaries composed typically of a mass-losing low-mass main-sequence secondary, an accreting white dwarf, and a luminous accretion disk. With a binary inclination of 83°, OY Carinae is one of only a handful of CVs in which the secondary eclipses the white dwarf. Details of OY Car’s two stars, the accretion disk, and the boundary layer between the accretion disk and the surface of the white dwarf are determined from eclipse observations in the optical, UV, EUV, and X-ray (Wood et al. 1989; Bruch et al. 1996; Horne et al. 1994; Naylor et al. 1988; Pratt et al. 1999).

One of the outstanding puzzles in the study of OY Car is the fact that in outburst the EUV flux is not eclipsed by the secondary (Naylor et al. 1988). The accretion disk wind is the most plausible source of this extended EUV emission, but it is not possible from the *EXOSAT* data to determine the *nature* of the emission. Possibilities include thermal emission and resonant scattering of boundary layer flux.

To determine the nature of the EUV emission of OY Car, we observed the source in superoutburst with the *Extreme Ultraviolet Explorer* (*EUVE*) satellite for 3 days in 1997 March. The resulting 80–180 Å spectrum, shown in Figure 1, contains broad (FWHM  $\sim 2000$  km s $^{-1}$ ) emission lines of O V–VI, Ne V–VI, Mg V–VI, and Fe VI–VIII. The pattern of emission lines is inconsistent with that of a thermal plasma, but we show below that it is reasonably well modeled by resonant scattering of boundary layer flux.

For the scattering model, we assume that the boundary layer radiates like a blackbody with a temperature  $T_{\text{bl}} = 100,000$  K and a luminosity  $L_{\text{bl}} = 4\pi R_{\text{wd}}^2 \sigma T_{\text{bl}}^4 = 4.4 \times 10^{34}$  erg s $^{-1}$  (where  $R_{\text{wd}} = 7.8 \times 10^8$  cm is the radius appropriate to a 0.7  $M_{\odot}$  white dwarf); the source is at a distance  $d = 85$  pc; and the intrinsic flux suffers photoelectric absorption by a column density  $N_{\text{H}} = 3 \times 10^{19}$  cm $^{-2}$ . For the scattering optical depth we assume  $\tau_{\lambda} = (\pi e^2 / m_e c) n_A \lambda_{ij} f_{ij} (g_i / \Sigma g_j) (dv/dr)^{-1}$  where the wind velocity gradient  $dv/dr = 3000$  km s $^{-1}$ /10 $^{10}$  cm;




Figure 1. Mean *EUVE* spectrum of OY Car. The spectrum is shown by the black histogram, the error by the grey histogram, the scattering model spectrum by the thick line. Line identifications are based on the model.

the density  $n = 10^{10} \text{ cm}^{-3}$  at a distance  $r = 10^{10} \text{ cm}$ ; the optical depth is distributed as a Gaussian with  $\text{FWHM} = 0.9 \text{ Å}$ ; the abundances  $A$  are as given by Allen (1973); and the wavelengths  $\lambda_{ij}$ , oscillator strengths  $f_{ij}$ , and statistical weights  $g_i$  are as given by Verner, Verner, & Ferland (1996). The resulting model shown in Figure 1 required only modest (factor of two) modifications of the abundances/ionization fractions to produce the match to the data.

Given the success of this model, it is natural to ask if radiation pressure alone is capable of driving the inferred mass-loss rate of  $\dot{M}_{\text{wind}} = 4\pi r^2 \mu m_{\text{H}} n v = 8 \times 10^{15} \text{ g s}^{-1}$ . It is not: for reasonable parameters, the total radiative force is  $\sim 200$  times that of the O VI  $\lambda 150$  line of  $L_{\text{O VI}}/c = 3 \times 10^{20} \text{ dynes}$ ; the resulting mass-loss rate is  $\dot{M}_{\text{wind}} = L_{\text{total}}/vc \sim 2 \times 10^{14} \text{ g s}^{-1}$ , an order of magnitude short of the value required.

C. W. M.'s contribution to this work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.

## REFERENCES

- Allen, C. W. 1973, *Astrophysical Quantities* (London: Athlone Press)
- Bruch, A., Beele, D., & Baptista, R. 1996, *A&A*, 306, 151
- Horne, K., Marsh, T. R., Cheng, F. H., Hubeny, I., & Lanz, T. 1994, *ApJ*, 426, 294
- Naylor, T., et al. 1988, *MNRAS*, 231, 237
- Pratt, G. W., Hassall, B. J. M., Naylor, T., & Wood, J. H. 1999, *MNRAS*, 307, 413
- Verner, D. A., Verner, E. M., & Ferland, G. J. 1996, *Atomic Data and Nuc. Data Tables*, 64, 1
- Wood, J. H., Horne, K., Berriman, G., & Wade, R. A. 1989, *ApJ*, 341, 974
- C. W. Mauche: Lawrence Livermore National Laboratory, L-43, 7000 East Avenue, Livermore, CA 94550 (mauche@cygnus.llnl.gov)
- J. C. Raymond: Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 01238 (jraymond@cfa.harvard.edu)